Prescription Opioids: What you need to know



Prescription opioids can be used to help relieve moderate-to-severe pain and are often prescribed following a surgery or injury, or for certain health conditions. These medications can be an important part of treatment but also come with serious risks. It is important to work with your health care provider to make sure you are getting the safest, most effective care.

What are the risks and side effects of opioid use?

Prescription opioids carry serious risks of addiction and overdose, especially with prolonged use. An opioid overdose, often marked by slowed breathing, can cause sudden death. The use of prescription opioids can have a number of side effects as well, even when taken as directed:

• Tolerance–meaning you might need to take more of a medication for the same pain relief

Physical dependence—meaning

when a medication is stopped

Increased sensitivity to pain

- Nausea, vomiting, and dry mouth
- Sleepiness and dizziness
- Confusion
- Depression
 - Low levels of testosterone that can result in lower sex drive, energy, and strength
 - Itching and sweating

Risks are greater with:

- History of drug misuse, substance use disorder, or overdose
- Mental health conditions (such as depression or anxiety)
- Sleep apnea

Constipation

- Older age (65 years or older)
- Pregnancy





Learn more www.cdc.gov/drugoverdose/prescribing/guideline.html

All of this content was taken in its entirety from the Centers for Disease Control's publication, Prescription Opioids: What You Need to Know

As many as 1 in 4 people*

receiving prescription opioids long term in a primary care setting struggles with addiction.

Avoid alcohol while taking prescription opioids. Also, unless specifically advised by your health care provider, medications to avoid include:

- Benzodiazepines (such as Xanax or Valium)
- **Muscle relaxants** (such as Soma or Flexeril)
- Hypnotics (such as Ambien or Lunesta)
- Other prescription opioids
- * Findings from one study



Know your options

Talk to your health care provider about ways to manage your pain that don't involve prescription opioids. Some of these options **may actually work better** and have fewer risks and side effects. Options may include:

- Pain relievers such as acetaminophen, ibuprofen, and naproxen
- Some medications that are also used for depression or seizures
- Physical therapy and exercise
- Cognitive behavioral therapy, a psychological, goal-directed approach, in which patients learn how to modify physical, behavioral, and emotional triggers of pain and stress.

If you are prescribed opioids for pain:

- Never take opioids in greater amounts or more often than prescribed.
- Follow up with your primary health care provider within <u>days</u>.
 - Work together to create a plan on how to manage your pain.
 - Talk about ways to help manage your pain that don't involve prescription opioids.
 - Talk about any and all concerns and side effects.
- Help prevent misuse and abuse.
 - Never sell or share prescription opioids.
 - Never use another person's prescription opioids.
- Store prescription opioids in a secure place and out of reach of others (this may include visitors, children, friends, and family).
- Safely dispose of unused prescription opioids: Find your community drug take-back program or your pharmacy mail-back program, or flush them down the toilet, following guidance from the Food and Drug Administration (www.fda.gov/Drugs/ResourcesForYou).
- Visit <u>www.cdc.gov/drugoverdose</u> to learn about the risks of opioid abuse and overdose.
- If you believe you may be struggling with addiction, tell your health care provider and ask for guidance or call SAMHSA's National Helpline at 1-800-662-HELP.



Be informed!

Make sure you know the name of your medication, how much and how often to take it, and its potential risks and side effects.

Learn more www.cdc.gov/drugoverdose/prescribing/guideline.html